



$\text{C}_2\text{H}_5\text{MgBr} + \text{CH}_3\text{COCl} \rightarrow \text{CH}_3\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{MgBr}$
 &fĩ ċ' ()* (+* ffl ' / ċ' (~ (Angew. Chem. Int. Ed. 2015, 54, . //01. //2(



&fĩ ċ' # 3 (~4 f5L'' fi
 6f5 4 ° 57
 -fĩ 5fi #. ċ /8. 2

9! f! 5L~ #/ ~! 1 1 1



F Racemic synthesis reported by G Hssinger in 1998.

F 33 steps longest linear sequence from thymoquinone.


> f! 1 1 ° = ° J (K (+> f! / t ° - (+4 H! 1 1 ° = ° i (+4 ° 5! t ° A (+* t / L ° 4 (Monatsh. Chem. 1994, 125, MM1. 8. 8 (M! t °, f! ~ 0
 ~~~~~ Z! = 5f! H O  
 > f! / t ° - (+4 H! 1 1 ° = ° i (+A P Z! f! C (+\* t / L ° 4 (+9 S = 1 ; t ° A (Angew. Chem. Int Ed. 1998, 37, ///21///E(

9°! fl 5L~ #/ ~ | 1 1 1

F Enantioselective synthesis achieved by Deslongchamps in 2003.

F

# NOV 17 11:41 AM



F Exceedingly acid-sensitive hemiketal moiety formed in the penultimate step.

F Intramolecular Diels-Alder cyclization with pyrone diene to set four of the final product's seven stereocenters.

F Vinylogous Mukaiyama aldol addition to combine two fragments containing all necessary carbon atoms.

D

X' °B fi; ~ #/ ~ 1 7fL


K\* 9YKt) \* fi'f'7i °fi; f'f'f'  
& fi' °) \* (+\* f'f' /' /' °) (- (Angew. Chem. Int Ed. 2015, 54, . //01. //2(  
i L'Z! tL, W(i (-5L'f'f'f' (-<fL; f'< (+X5f' /' 1L, [ (+9f'f' fL, \* (+% ~ ' \5f' f'Z) 6fL9.

# X' °B fi; ~ #/ ~ ~ t 7fl



6K > Y ZT f5, ~; flL 7° 7#Zi, flt +K\* 9YKt 7t)\* fi "fl ~ 7t °fl; fl flt +K: 9t > Y N, N-; flL 7° 7#Z) t ~ #Zi flt  
& fi ~ 0)\* (+\* fl) /t ~ (- (Angew. Chem. Int. Ed. 2015, 54, . //01. //2(  
Xt ~ t ~ > +%fi t flfi i (~\* (J. Org. Chem. 2003, 68, MUQ1MEO(

# X' °B fi; ~ #/ ~ 1 7fl



F Diastereomers 7 and 8 separable by column chromatography.

F Only decarboxylated products obtained when attempted with triflate 5.


F Significant decarboxylation also observed in more polar solvents (DMF, acetonitrile), and at higher temperatures (120 °C).

F Cyclization reaction was markedly sluggish at 80 °C.

E



X' °B fi; ~ #/ ~ 1 7fl



M

& fi ~ ()\* (+\* fl) ' /t ~ (- (Angew. Chem. Int. Ed. 2015, 54, . //01. //2(



\* 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100



1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

\* 1 ž f/f, t



./

\* f#1 ° (ř6fl< (Angew. Chem. Int. Ed. 2006, 45, Q M01Q M0X)





\* 1 ž ~ f/x fl, 7

\* 1 ž ~ f i / f l , 7

.D

& fi 2 0\* (+\* ffl) ' / 2 2 (-(-

\* 1 ž ~ fi/fl, t



FA hemiketal formation mechanism involving intramolecular attack of a zincate on the lactone carbonyl is also plausible:

